

Towson University Interlibrary Loan



ILLiad TN: 36462

Borrower: GDC

Lending String: VXF,*TSC,OKX,UPM,UPM

Patron: ;dept; ;type; Amith, Jonathan

Journal Title: Phytologia.

Volume: 72(3);231-252. **Issue:** 3

Month/Year: 1992**Pages:** 231-52

Article Author:

Article Title: Nesom, G. L.; New species and taxonomic evaluations of Mexican *Castilleja* (Scrophulariaceae).

Imprint: Huntsville, TX [etc.] Warner Herbarium [

ILL Number: 19552518



Call #:

Location: bound

ARIEL

Charge

Maxcost: \$25IFM

Shipping Address:

Gettysburg College

Library

Interlibrary Loan

Gettysburg, PA 17325-1493

Fax: 717-337-7001

Ariel: 138.234.152.5

_____. 1971. A taxonomic revision of the *Castilleja viscidula* group. Mem. New York Bot. Gard. 21:1-63.

_____. 1976. Four new species of Mexican *Castilleja* (subgenus *Castilleja*, Scrophulariaceae) and their relatives. Brittonia 28:195-208.

_____. 1984. *Castilleja*. *Intermountain Flora* 4:476-496.

Johnston, M.C. 1990. *The Vascular Plants of Texas*. A list, updating the *Manual of the Vascular Plants of Texas*, 2nd ed. Published by the author, Austin, Texas.

Lockwood, M. 1991. Systematic evaluation of five species of *Castilleja* (Scrophulariaceae) in trans-Pecos, Texas. M.S. Thesis, Sul Ross State University, Alpine, Texas.

Lockwood, M. & M. Forstner. 1991. In IOPB chromosome data 3. Intern. Org. Pl. Biosys. News. 17:10.

Nesom, G.L. 1992. New species and taxonomic evaluations of *Castilleja* from México. Phytologia 72:231-252.

Pennell, F.W. 1940. Scrophulariaceae of trans-Pecos Texas. Proc. Acad. Nat. Sci. Philad. 92:289-308.

Standley, P.C. 1909. More southwestern castillejas. Muhlenbergia 5:81-87.

Ward, D.E. 1983. Chromosome counts from New Mexico and southern Colorado. Phytologia 54:302-309.

Ward, D.E. 1984. Chromosome counts from New Mexico and Mexico. Phytologia 56:55-60.

Warnock, B.H. 1970. *Wildflowers of the Big Bend Country, Texas*. Sul Ross State Univ., Alpine, Texas.

_____. 1977. *Wildflowers of the Davis Mountains and Marathon Basin, Texas*. Sul Ross State Univ., Alpine, Texas.

Wootton, E.O. & P.C. Standley. 1915. *Flora of New Mexico*. Contr. U.S. National Herbarium, Vol. 19.

NEW SPECIES AND TAXONOMIC EVALUATIONS OF MEXICAN *CASTILLEJA* (SCROPHULARIACEAE)

Guy L. Nesom

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

ABSTRACT

Four new species of *Castilleja* are described from México. *Castilleja jiquilpana* from Michoacán is most closely related to *C. scorzoneraefolia*. *Castilleja durangensis* from Durango is most closely related to *C. aspera*, and *C. galehintoniae* from Nuevo León is most closely related to *C. lanata*. *Castilleja dendridion* from Oaxaca has much more uncertain evolutionary affinities. The identities and taxonomic status of the accepted species *C. scorzoneraefolia*, *C. falcata*, *C. rigida*, *C. nervata*, *C. bella*, and *C. aspera*, as well as taxa relegated to synonymy, are evaluated.

KEY WORDS: *Castilleja*, Scrophulariaceae, México

In the course of providing general identifications of recent collections of *Castilleja* from northern México, as well as curating LL, TEX collections of the genus, a number of taxonomic problems and undescribed taxa have come to light. A review of the genus in México and Central America (Nesom unpublished) accounts for 52 species in mainland México, an additional thirteen in Baja California (only one of these occurs on the mainland), and nine in Central America (six endemic there). Eastwood's study (1909) of Mexican and Central American *Castilleja* recognized 54 species, seventeen originally described by her. Brandegee (1914) described an additional species from mainland México, Standley & Steyermark (1944) one from Guatemala, Standley (1936; 1940) three species from mainland México and one (1938) from Costa Rica, the latter including three species later proposed by Pennell, Crosswhite (1970) one, Rzedowski (1975) one, Holmgren four from México (1976) and four from Costa Rica and Panamá (1978), and Moran (*in* Levin & Moran 1989) one from México. Breedlove & Heckard (1970) placed a new species from Sinaloa in the monotypic genus *Gentrya*, which has recently been positioned within

Castilleja (Chuang & Heckard 1991). Six other species have been described since 1909 from Baja California or from southern California and are known to occur in Baja California. The present study adds four new species of *Castilleja* from mainland Mexico: a companion paper (Nesom 1992) adds another that is primarily centered in Texas but that also occurs in Coahuila.

Where species have been problematic in their past circumscriptions, their identity and geographical distribution are documented by specimen citations in the present study. All taxa are mapped, based primarily on specimens from GH, MO, NY, and LL, TEX.


I. The identity of *Castilleja scorzoneraefolia* Kunth

Castilleja scorzoneraefolia Kunth, *Nov. Gen. & Sp.* 2 [folio]:267. 1818; 2 [quarto]:331, tab. 165. 1820. TYPE: MEXICO. Michoacán: "Ario Pazcuaro" (as on the specimen), [Sep 1803], *Humboldt & Bonpland s.n.* (HOLOTYPE: P fide!). The protologue: "Crescit in rupestribus Novae Hispaniae prope Pazcuaro et Ario, alt 1100 hex. 4 Floret Augusto."

Castilleja lithospermoides Kunth, *Nov. Gen. & Sp.* 2 [folio]:266. 1818; 2 [quarto]:331, tab. 164. 1820. TYPE: MEXICO. Hidalgo: "Moran" (as on the specimen), [May-Jun 1803], *Humboldt & Bonpland s.n.* (P fide!). The protologue: "Crescit in temperatis Novae Hispaniae prope Real del Monte et Moran, et in Regno Quitense prope Chillo, alt 1350 hex. 4 Floret Mayo-Junio." The sheet of *C. lithospermoides* has 3 branches: the one on the left is typical *C. scorzoneraefolia*, the identity of the other two is more equivocal, and it is possible that they might even represent what is identified in the current study as *C. nervata* Eastwood. If so, however, it is not likely that they originated from central Hidalgo, which is north of the range of *C. nervata* (Map 1). Further, the original illustration of *C. lithospermoides* (tab. 164) shows a plant nearly identical to that pictured for *C. scorzoneraefolia* (tab. 165), the leaves with expanded and distinctly clasping bases and corollas exerted from calyces.

Castilleja glandulosa Greenm., *Proc. Amer. Acad. Arts* 41:247. 1905. TYPE: MEXICO. Mexico: hills near Lecheria Station, 2200 m, 2 Jul 1904, *C.G. Pringle* 10,000 (HOLOTYPE: GH; Isotypes: LL, MO!).

Perennials usually from a taproot. Stems herbaceous, 10-30(40) cm tall, erect, straight, commonly with short, eglandular, deflexed hairs, stipitate glandular hairs, and much longer, stiffly spreading, eglandular hairs; lower leaf surfaces moderately to densely hirsute with stiff, straight, eglandular hairs,



Map 1. Distribution of *Castilleja nervata*, *C. scorzoneraefolia*, *C. jiquilpana*, and *C. dendridion*. Arrows show the locations of type collections.

sometimes also with glandular hairs, the margins commonly ciliate with much longer hairs. Leaves 2-7 cm long, narrowly triangular, entire, 3 veined, the veins sometimes prominently raised, sessile, distinctly clasping. Floral bracts 24-30 mm long, oblong lanceolate to obovate, 3 veined, 5-12 mm wide, entire, the upper red tipped. Calyx red and glandular at the apex, 19-32 mm long, from the base is somewhat constricted at ca. midlength, then broadening again toward the apex, equally divided, the primary lobes 7-12 mm long, the secondary lobes rounded and shallow, 1-3 mm long, sometimes completely absent. Corolla 21-37 mm long, the lower lip of 3(-5), thick green teeth 1-2 mm long, the galea 7-12 mm long, 30%-34% as long as the corolla, usually exerted from the calyx 3-9 mm, rarely included. Chromosome numbers, $n=12$ and $n=24$, both unpublished counts by Chuang, according to label data.

Widespread in temperate areas of southcentral to northeastern México (Map 1); (950-1250-3590(-4100) m; Apr-Sep).

Castilleja scorzoneraefolia is one of the most common and widespread species of México. It is somewhat variable in features of vestiture, calyx morphology, and degree of corolla exertion, but for the most part, identifications of this species are relatively consistent. Possible intermediacy between *C. scorzoneraefolia* and *C. rigida* Eastwood and *C. nervata* is discussed below. Numerous collections from westcentral Zacatecas and adjacent Durango appear to represent a disjunct, northwesternmost segment (Map 1) of *C. scorzoneraefolia* and may prove to deserve taxonomic recognition. These plants consistently produce highly glandular stems and leaves, leaves with sinuate margins, prominent axillary clusters of leaves, and calyces with red and white longitudinal stripes. Such features, however, also occur in plants scattered through the range of the species.

Eastwood (1909) distinguished *Castilleja scorzoneraefolia* from close relatives by its lower corolla lip of five teeth, three primary ones and two smaller ones in the sinuses of the others. I also have observed this feature, but it is not constant within the species. A similar lower corolla lip also occurs sporadically in plants of *C. rigida* (see below).

Plants with 3 lobed bracts and leaves, but otherwise mostly similar to *C. scorzoneraefolia*, occur in the high elevation regions of Veracruz (Orizaba and Perote, Map 1), where they have been named as *C. falcata* Eastwood. These apparently are annual or short lived perennials and are erect and single stemmed from the base. They appear to intergrade with typical *C. scorzoneraefolia*, but no other plants of the latter over the range of the species produce this morphology and *C. falcata* is tentatively maintained here as a taxon deserving recognition at least at some rank.

Castilleja falcata Eastwood, Proc. Amer. Acad. Arts 44:575. 1909. TYPE: MEXICO. Puebla: Mount Orizaba, 3660 m, 14 Aug 1901, C.G. Pringle

8560 (HOLOTYPE: GH; Isotype: MOJ).

In contrast to the somewhat ambiguous distinction of *Castilleja falcata* from *C. scorzoneraefolia*, another closely related but previously unrecognized population system appears to be clearly distinct as a species and is formally described here.

Castilleja jiquipana Nesom, sp. nov. TYPE: MEXICO. Michoacán, 1 mi

W of El Fresno, 6 mi from jet of Hwy 15 on Hwy 110, 6000 ft, 7 Jul 1966, E. Molsseed 441 (HOLOTYPE: LL; Isotypes: MOJ, UC).

Castilleja scorzoneraefoliae Kuntz similis sed duratione brevior, caulium vestimento sparsim piloso glanduloso, foliorum lobbis linearibus, et calycibus flavifasciatis differt.

Plants apparently annual, from a short, slender taproot. Stems 22-38 cm tall, eglandular, very sparsely pilose with vitreous, spreading hairs mostly 0.5-1.0 mm long and a sparse understorey of much smaller, loose, often somewhat deflexed hairs. Leaves subclasping, lanceolate, 2-5 cm long, 2-7 mm wide (at midleaf), 3 veined with the midvein raised, at least the lower leaves with 1-3 pairs of filiform lobes from the upper half of the broad central portion of the lamina, upper leaves entire to lobed. Mature inflorescence/infructescence 8-15 mm long; floral bracts lanceolate, the lowermost sometimes lobed like the leaves, 17-25 mm long, 4-6 mm wide, the upper third red and minutely glandular. Calyx red tipped and glandular at the apex, 15-17 mm long, from the base is somewhat constricted at ca. midlength, then broadening again toward the apex, the primary lobes 6-8 mm long, nearly equal in length, rounded at the apex with barely, if at all developed secondary lobes, or shallowly notched with small but distinct secondary lobes, red on the upper fourth, green below, a yellow band ca. 1 mm wide between the red and green portions. Corolla exerted from the calyx 1-4 mm, the lower lip of 3, thick, green teeth ca. 1 mm long, the galea 6-8 mm long, 33%-45% as long as the corolla, glandular and sparsely pilose. Chromosome number, $n=12$ (see voucher below).

Northwest Michoacán (Map 1); wet, grassy fields and roadsides, 2050-2310 m, Jun-Aug.

Additional collections examined: MEXICO. Michoacán: Ca. 22 km W of Jiquipán on Rta. 110, 3 Aug 1966, Cruden 1149, voucher for chromosome count of $n=12$, as *Castilleja glandulosa* [Heckard 1968] (TEX); 12-13 km W of Jiquipán on Rta. 110, 1 Jul 1968, Cruden 1322 (NY).

Castilleja jiquipana apparently is a narrow endemic (Map 1) in a distinctive habitat. All three collections studied were made within about 30 kilometers of each other in the area of Jiquipán, Michoacán. The plants are similar to *C. scorzoneraefolia* particularly in their subclasping leaves and features of

the calyx. The new species is morphologically distinct in its apparently annual duration (judging from the short, very slender taproot), yellow banded calyces, eglandular and sparsely pilose stems, and deeply divided leaves with linear lobes. Scattered plants of *C. scorzoneraefolia* throughout its range may show some of these features, but the only other plants within the *C. scorzoneraefolia* group with similarly divided leaves and bracts are those of *C. falcata*, *C. moranensis* Kunth, and *C. bella* Standley (see discussion below of the latter two).

Castilleja jiguiripana also is at least superficially similar to *C. saltenensis* Eastwood, which is endemic to southeastern Durango. Plants of the latter, however, are shorter with floral bracts that are most often lobed (vs. entire), a more villous stem vestiture of hairs with a strong tendency to be produced in vertical lines, and calyces that lack a yellow band beneath the distal red portion.


II. The identity of *Castilleja rigida* and *C. nervata*

Castilleja rigida Eastwood, Proc. Amer. Acad. Arts 44:575. 1909. TYPE: MEXICO. Chihuahua: hills near Chihuahua, 16 Apr 1885, C.G. Pringle 188 (HOLOTYPE: GH!).

Castilleja latebracteata Pennell, Proc. Acad. Nat. Sci. Philad. 73:533. 1921. TYPE: UNITED STATES. Texas: Val Verde Co., High Bridge [Southern Pacific Railroad] of the Pecos, 1000 ft, 27-28 Apr 1903, Dr. H.A. Pilsbry s.n. (HOLOTYPE: PH!).

Stems woody, (15-)14-55 cm tall, commonly slightly zig-zag at the nodes. Stem and leaf surfaces softly pilose to villous with eglandular, vitreous, loosely spreading, and somewhat crinkled hairs, mixed with gland tipped hairs. Leaves obovate, sessile, not clasping, without prominently raised veins. Floral bracts entire, 3 veined, 8-16 mm wide. Calyx red tipped and glandular at the apex, 20-32(-36) mm long, from the base is somewhat constricted at ca. midlength, then broadening again toward the apex, the primary lobes lobes (5-)7-12 mm long, nearly equal in length, with rounded, barely developed secondary lobes. Corolla conspicuously exerted from the calyx, (24-)28-36(-42) mm long, the lower lip of 3(-5), thick, green teeth 0.5-1.5 mm long, the galea (8-)10-14 mm long, 33%-45% as long as the corolla. Chromosome number unknown.

Chihuahuan Desert region of Chihuahua, Durango, Zacatecas, Coahuila, and Nuevo León, México, also trans-Pecos Texas (Map 2); matofrall, sometimes into mesquite or oak woods, limestone substrates, except in the igneous Santa Rosa Mountains southwest of Muzquiz; (500-)1100-2800 m; (Mar-)Apr-Oct (-Feb).



Map 2. Distribution of *Castilleja rigida*, *C. aspera*, and *C. durangensis*. Arrows show the locations of type collections.

Castilleja rigida is the earliest and correct name for the plants of trans-Pecos Texas and northern México that previously (Holmgren 1970) have been identified as *C. latebracteata* Pennell. The species is almost completely confined to the Chihuahuan Desert Region and occurs relatively abundantly in the Mexican states of Durango, Zacatecas, Chihuahua, Coahuila, and Nuevo León, as well as in southern Texas (Map 2; also see Nesom 1992). The plants are easily identified by their morphology and geographic range and I have not cited specimens.

In its calyx and corolla morphology, *Castilleja rigida* is similar to *C. scorzoneraefolia*, and the two species are almost certainly closely related. There is some indication that gene flow occurs between them where they are sympatric in Coahuila. For example, in the Sierra de la Gavia (35 mi south of Monclova), Henrickson has collected plants of *C. rigida* (oak woods at 4500 ft, 117776, NY-2 sheets) with tall, slightly zig-zag stems and nonclaspng leaves but atypical in their very narrow floral bracts, in close proximity to *C. scorzoneraefolia* (oak chaparral at 5900 ft, 11808, NY-2 sheets) with clasping leaves but tall, distinctly woody stems and wide floral bracts. Similar examples can also be found in the region of Muzquiz, Coahuila, at the northernmost extension of the range of *C. scorzoneraefolia*.

Castilleja rigida also has been confused with *C. nervata* Eastwood (in specimen annotations, by my own and those of others, the two species considered together as *C. nervata*). Plants of both species are strongly perennial with woody, slightly zig-zag stems and non-claspng leaves, but they are distinct from each other in both geography (Maps 1 and 2) and a number of morphological features. Both *C. rigida* and *C. nervata* are closely related to *C. scorzoneraefolia*. The three species are distinguished by features in the following couplet.

1. Plants usually with a distinct taproot; stems straight, herbaceous from the base; leaf bases expanded, clasping to subclasping; calyx red tipped; corollas exerted from the calyx. *C. scorzoneraefolia*
1. Plants usually with woody roots but not distinctly taprooted; stems slightly zig-zag at the nodes, basally herbaceous to woody; leaves attenuate to a nonclasping base; calyx red tipped or completely green; corollas included within the calyx or exerted. (2)
2. Corollas (24-)28-36(-42) cm long, exerted from the calyx, the lower lip of linear-attenuate teeth; calyx red tipped; leaves with primary abaxial foliar veins not prominently raised; stem and leaf surfaces softly pilose to villous with eglandular, loosely spreading, and some what crinkled hairs, mixed with gland tipped hairs, the leaf margins essentially ciliate. *C. rigida*

2. Corollas 15-24 cm long, completely included within the calyx, the lower lip of deltate or triangular teeth; calyx completely green; primary abaxial foliar veins prominently raised; stems commonly with short, eglandular, deflexed hairs, stipitate glandular hairs, and much longer, stiffly spreading, eglandular hairs, the lower leaf surfaces moderately to densely hirsute with stiff, straight, eglandular hairs, sometimes also with glandular hairs, the margins commonly ciliate with much longer eglandular hairs. *C. nervata*

Castilleja nervata Eastwood, Proc. Amer. Acad. Arts 44:574, 1909. [22 May] TYPE: MEXICO. Chihuahua: vicinity of Madera, 2250 m, May-Jun 1908, E. Palmer 274 (HOLOTYPE: GH; Isotype: MO!).

Castilleja angustifolia Mart. & Gal., Bull. Acad. Roy. Sci. Brux. 12(2):29, 1845. TYPE: not seen. Not *Castilleja angustifolia* G. Don, Gen. Syst. Gard. Bot. 4:616, 1838, from North America; not *Castilleja angustifolia* (Nutt.) A. Gray in Torrey, Bot. U.S. & Mex. Bound. Surv. 2(1):118, 1838, from North America. The identity of *C. angustifolia* Mart. & Gal. is accepted here *vide* Eastwood (1909), who noted that it is the same species as *C. lithospermoides* (= *C. nervata* in the sense of the present treatment, see comments below). The name upon which Gray's combination was based, *Euchroma angustifolia* Nutt., almost certainly represents some other species, perhaps *C. angustifolia* (Nutt.) G. Don, rather than the specimen Gray referred to, which was apparently at hand (= *Castilleja integra* A. Gray, a Wright collection from Texas).

Castilleja cryptandra Eastwood, Proc. Amer. Acad. Arts 44:578, 1909. TYPE: MEXICO. Colima: Cuchilla, NE side of Volcan Colima, 22 Jul 1905, P. Goldsmith 76 (HOLOTYPE: GH; Isotype: MO!).

Castilleja cruenta Standley, Muhlenbergia 5:82, 1909 [7 July]. TYPE: UNITED STATES. Arizona: Cochise Co., Chiricahua Mountains, rocky spur N of Wilgus Ranch, rhyolite, 6000 ft, 2 Sep 1907, J. C. Blumer 2193 (HOLOTYPE: NY! ex NMC). The identity of these plants with *Castilleja nervata* has also been recognized by N. Holmgren, as evidenced by his 1983 annotation of the type specimen.

Stems woody, (15-)30-100 cm tall, commonly slightly zig-zag at the nodes. Stems commonly with short, eglandular, deflexed hairs, stipitate glandular hairs, and much longer, stiffly spreading, eglandular hairs; lower leaf surfaces moderately to densely hirsute with stiff, straight, eglandular hairs, sometimes also with glandular hairs, the margins commonly ciliate with much longer

hairs. Leaves obovate to linear lanceolate, sessile, not clasping, the 3 major veins prominently raised on the lower surface. Floral bracts entire, 3 veined, 6-15 mm wide. Calyx green, 15-24 mm long, tubular, the primary lobes 8-14 mm long, nearly equal in length, with rounded, barely developed secondary lobes. Corolla completely included within the calyx, 15-24 mm long, the lower lip of 3, green, thickened; linear-attenuate teeth 2.0-2.5 mm long, the galea 5-8 mm long, ca. 33% as long as the corolla. Chromosome numbers, $n=12$, $n=24$ (see vouchers below).

Southern Arizona in the United States, and in the following Mexican states: Sonora, Chihuahua, Distrito Federal, Sinaloa, Durango, Nayarit, Jalisco, Colima, México, Michoacán, Guerrero, Morelos, Puebla, Veracruz, and Oaxaca (Map 1); rocky slopes or openings, oak to pine-oak or pine woods, rarely in pine-fir; 750-2300(-2600) m. (Apr.-Jul-Oct.-Nov.). Eastwood (1909) noted that this species (as *Castilleja lithospermoides*, see comments below) occurs in South America, but I have confirmed its presence only in México.

Representative collections examined: MEXICO. Chihuahua: Mpio. Batopilas, N of Quitare on La Bufa-Creel road, 31 Jul 1977, *Bye 7782* (GH); Mesa de Arroyo Seco, 25-30 km SW of Minaca, 16-17 Sep 1994, *Pennell 18846* (GH, NY). Distrito Federal: Lomas, Sep 1930, *Lyonnét 749* (MO). Durango: Along trail from Pueblo Nuevo to Cueva, 30-31 Aug 1934, *Pennell 18487* (GH). Guerrero: SW of Xochipala, ca. 40.5 km SW of K236 on Rta. 95, 20 Jul 1969, *Cruden 1628* (NY). Jalisco: Ca. 17.5 km S of Autlán on Route 80, 2 Aug 1969, *Cruden 1672* (NY); hills near Guadaluajara, 13 May 1901, *Pringle 9461* (GH, NY). México: 10 mi N of México City near Atzacapotzotco, 1-15 Jul 1937, *Happ 109* (MO). Michoacán: Cerro Tancitaro, 19 Aug 1940, *Leauenworth 664* (NY, with *Castilleja scorzoneraefolia*); hills near Patzcuaro, 30 Jul 1982, *Pringle 4168* (GH, MO, NY). Morelos: Cuautla, Jul 1930, *Lyonnét 743* (NY); La Herradura near Cuernavaca, 14 May 1938, *Williams 8048* (MO). Nayarit: Cerro de San Juan, SW of Tepic, 18 Aug 1935, *Pennell 19754* (GH, NY); ca. 19 km S of Tepic, 27 Jun 1968, *Cruden 1913* (NY). Oaxaca: Dpt. Mixe, between San Isabel and San Juan Mazatlán, [no date], *Lapp 37* (NY); Dept. Etla, 17 km NE of San Gabriel Etla, 13 May 1985, *Lopez G. 258* (NY). Puebla (?): Orizaba, Engenio, Sierra de Cruz, Aug 1853, *Müller s.n.* (NY). Sinaloa: Cerro de la Sandia, NE of Panuco, 29-30 Aug 1935, *Pennell 20050* (GH). Sonora: Puerta de Pintos, 14 Oct 1890, *Hartman 150* (GH); Bakachaka, Río Mayo, 5 Jul 1935, *Gentry 1464* (GH, MO); ridge S of Arroyo Gochico, E of San Bernardo, 5-9 Aug 1935, *Pennell 19531* (GH, NY). Veracruz: Mt. Orizaba, Magayas to Lomogrande, 26 Apr 1938, *Balls 4947* (GH).

UNITED STATES. Arizona: Santa Cruz Co., Santa Rita Mountains, 7000 ft, 25 Jul 1884, *Pringle 8175* (NY) and *8174* (NY). Eastwood (1909) recognized *Castilleja nervata* only from Chihuahua. The remainder of the species (as treated here) was identified by her as *C. lithospermoides*, although she noted that the two taxa were similar. The type of *C.*

lithospermoides, however, was collected in Hidalgo (north of the known range of *C. nervata*), and the type specimen (fiche) as well as the original illustration show clasping leaves and exerted corollas characteristic of *C. scorzoneraefolia*. Jiménez (1985) clearly used *C. lithospermoides* as the name for what is identified in the current study as *C. scorzoneraefolia*; the plants he identified as *C. scorzoneraefolia*, with acute calyx lobes and large anthers, are some other species.

The type of *Castilleja cruenta* (Chiricahua Mountains, Arizona), is typical of *C. nervata*; the plants from the Santa Rita Mountains, Arizona (*Pringle 8174* and *8175*), are atypical in their shallowly toothed floral bracts but otherwise so similar to *C. nervata* that they must be referred to it.

Castilleja nervata and *C. scorzoneraefolia* have closely contiguous geographic distributions (Map 1), and apparent intermediates may be found along the area where their ranges abut. The two species are sympatric in the area of Edo. México, Morelos, the north half of Oaxaca, and Puebla, the area of Volcán de Colima along the Jalisco-Colima border, the area of Mt. Tancitaro, Michoacán, and in southwestern Durango. Within *C. scorzoneraefolia*, scattered through its range, aspects of variability support an hypothesis of close relationship to *C. nervata*. Although the corollas are usually well exerted in *C. scorzoneraefolia*, they are sometimes nearly included, and although the vestiture is mostly of loose hairs, they may be distinctly hispid.

III. The identity of *Castilleja bella*

Castilleja bella has previously been known to occur on Cerro Potosí, Nuevo León (the type locality) and Cerro Peña Nevada (along the Nuevo León-Tamaulipas border) (McDonald 1990), and its presence is here recognized on the high peaks east of Saltillo, Coahuila (Sierra Coahuilán and Sierra La Marta). The extension of its known range to the latter area makes its distribution more typical of the majority of species of the alpine-subalpine flora of northeastern México (McDonald in press).

Castilleja bella Standley, Field Mus. Pub. Bot. 22:106. 1940. TYPE. MEXICO. Nuevo León: Mpio. Galeana, peak of Cerro Potosí, abundant in meadows above and below timber line, 21 Jul 1935, *C.H. Mueller 2949* (HOLOTYPE: F; Isotype: MO!).

Caespitose perennials from a thick, straight taproot. Stems herbaceous, 2-6(-30) cm tall, stems and leaves very sparsely pilose with loose, vitreous hairs mostly 0.8-1.6 mm long, eglandular. Leaves 2-5 cm long, narrowly triangular, 3 veined, sessile, distinctly clasping, with 1-2 pairs of narrow lobes or the taller plants with at least the lower leaves entire. Floral bracts 23-28 mm

long, oblong lanceolate to obovate, 3 veined, 3-5 mm wide, with 1-2 pairs of narrow lobes, the upper 2/3 of each lobe red tipped. Calyx red and glandular at the apex, 28-36 mm long, equally divided, the primary lobes 6-10 mm long, the secondary lobes rounded and shallow, 1-2 mm long, sometimes completely absent. Corolla 28-38 mm long, the lower lip of 3, fleshy, green teeth 0.5-2.0 mm long, the galea 6-9 mm long, 25%-33% as long as the corolla, exerted from the calyx 1-6 mm. Chromosome number unknown.

High sierra of southeastern Coahuila (Sierra La Marta and Sierra Coahuilón), northcentral Nuevo León (Cerro Potosí), and southern Nuevo León and adjacent Tamaulipas (Cerro Peña Nevada) (Map 4), alpine and subalpine zones, 3000-3700 m; May-Jul-Nov.; at the crests of Sierra La Marta and Sierra Coahuilón among *Pinus culminicola* Andresen & Beaman and shrubs, pine-fir meadows, pine-oak woods, 3016-3600 m; on Cerro Potosí in rocky, alpine meadows and edge of krumholz near the peak, into pine meadows below, 3350-3700 m; on Peña Nevada in the area of the highest peak, 3400-3600 m.

Additional collections examined: MEXICO. Coahuila: Mpio. Arteaga: Sierra Coahuilón, 17 Jun 1991, *Hinton et al.* 20998 (TEX); Sierra La Marta, 17 May 1981, *Poole* 2916 (TEX); Sierra La Marta, first peak E of Cerro Morro, 20 Jul 1985, *McDonald* 1685 (TEX). Nuevo León: Mpio. Dr. Arroyo, Sierra de Peña Nevada: north of Picacho de San Onofre, burned zone on W side, with dominant *Ceanothus* and *Arctostaphylos*, ca. 3400 m, 30 Nov 1984, *McDonald* & *Gómez* 1292 (TEX); Picacho de San Onofre, E side, ca. 3600 m, 5 Jul 1985, *McDonald* 1661 (TEX); Mpio. Galeana, Cerro Potosí: NE summit, 13 Sep 1960, *Beaman* 4456 (GH); near summit, 27 Aug 1987, *Bogler* & *Atkins* 162 (TEX); summit, 18 May 1982, *Dorr* 2277 (TEX); top, 28 Oct 1982, *Grimes* 2971 (TEX); Cerro Potosí ascent, 3350 m, 2 May 1969, *Hinton et al.* 17012 (TEX); [near summit], 21 Jun 1969, *Hinton et al.* 17109 (TEX); summit, 25 May 1969, *Hinton* 17049 (TEX); top, 23 Aug 1984, *Lawn* 4781 (TEX); alpine zone, 26 Oct 1984, *McDonald* & *Gómez* 1260 (TEX); alpine zone, 26 Jul 1985, *McDonald* 1788 (TEX); ascent of Sierra Potosí by the N hogback, abundant on the meadow at the peak and common in the forest immediately beneath, 26 Jul 1934, *Mueller* 1234 (GH, TEX); ascent of Sierra Potosí by the north hogback, common in all parts of the pine forest, 26 Jul 1934, *Mueller* & *Mueller* 1239 (GH, TEX); peak, 18 Jul 1938, *Unu. Illinois Exped.* 935 (GH, MO); summit, 20 Oct 1979, *Wormock* 2013 (TEX); summit, 24 Jul 1977, *Wells* & *Nesom* 295 (TEX). Tamaulipas: Mpio. Miquihuana, Cerro Peña Nevada, 1 Jun 1974, *Patterson* 1514 (TEX).

In the original description of *Castilleja bella*, Standley (1940, p. 106) quoted comments by the collector of the paratype regarding variation in the species: "A very conspicuous plant, abundant on the treeless peak, and extending down (equally abundant) into the pine forest and in grassy openings for over 300 meters. In exposed places only 2.5-5 cm high. In the shelter of the timber at times reaching a height of 12 cm, but even in such places most of the plants are

low." The same variation in habit noted by Standley is confirmed by study of the numerous herbarium specimens by which this species is now represented. The plants are mostly acaulescent (2-6 cm tall) in alpine zones, but even on the peak (and near it) of Cerro Potosí, among acaulescent plants are individuals with stems commonly to 12 cm tall and ranging up to 30 cm (e.g., *Hinton* 17109, *Mueller* 1234, *McDonald* 1788, *Bogler* & *Atkins* 162). Plants with the typical, acaulescent habit occur on Cerro Potosí and Peña Nevada, but so far only taller plants have been collected from Coahuila.

Castilleja bella differs from *C. scorzoneraefolia* in its high elevation habitats, caespitose habit, with numerous branches arising immediately from a thick taproot, floral bracts (and sometimes upper leaves) with 1-2 pairs of narrowly lanceolate lobes, and sparsely villous, glandular stems. The caespitose habit, however, as well as the reduced vestiture and similar taproots, are also found in some plants of *C. scorzoneraefolia* from Nuevo León and Coahuila, particularly at the upper range of elevation of that species. The primary distinction between *C. scorzoneraefolia* and the taller plants of *C. bella* is the lobing of the floral bracts and upper leaves. In Coahuila and Nuevo León, *C. scorzoneraefolia* occurs at 970-2700 meters elevation, with a few plants reaching as far up as 3400 meters. The lowermost elevation known for plants of *C. bella* is 3000 meters, but most occur above 3300 meters.

The similarities between *Castilleja bella* and high elevation *C. scorzoneraefolia* suggest that some gene flow may be occurring between the two taxa. In a few collections of relatively taller plants, some plants have lobed bracts, while others have entire bracts (e.g., *Mueller* & *Mueller* 1239-TEX lobed but *Mueller* & *Mueller* 1239-GH entire; *McDonald* 1695 and *Hinton* 17109, most plants with lobed bracts, few with entire). Somewhat arbitrarily, I have identified all plants from these populations as *C. bella*. On Peña Nevada, however, where *C. scorzoneraefolia* is common at lower elevations and extends upward to at least 3400 meters, the difference between it and *C. bella* is marked. No plants there of *C. scorzoneraefolia* produce lobed leaves or bracts and most of them are single- or few-stemmed from the base.

Castilleja bella is very similar to *C. moranensis* Kunth (= *C. pringlei* Fern. and *C. schaffneri* Hemsley), which is also primarily caespitose but shows the same variability in height as *C. bella*. *Castilleja moranensis* occurs in the states of Hidalgo, México, Morelos, Puebla, and Veracruz, where it is found in a variety of habitats in elevations ranging 2400-3700 (4000) meters. It is not clear whether *C. bella* and *C. moranensis* are sister species, with the caespitose habit and lobed floral bracts inherited from a common ancestor, or whether each species may have been independently derived from *C. scorzoneraefolia*. The two can be distinguished by the contrasts in the following couplet.

1. Calyx cinereous, the veins and lobe margins densely ciliate with stiffly

spreading, white hairs; leaves mostly 1-3 mm wide at midleaf (below the divergence of the lobes). *C. moranensis*

1. Calyx green, the veins and lobe margins sparsely pilose with loosely spreading, vitreous hairs; leaves mostly 4-6 mm wide at midleaf. *C. bella*

IV. The identity of *Castilleja aspera*

Castilleja aspera Eastwood, Proc. Amer. Acad. Arts 44:580. 1909. TYPE: MEXICO. Chihuahua: Sierra Madre near Colonia Garcia, 2287 m, 3 Jun 1899, C.H.T. Townsend & C.M. Barber 449 (HOLOTYPE: GH!).

Castilleja nelsonii Eastwood, Proc. Amer. Acad. Arts 44:579. 1909. LECTOTYPE (designated here): MEXICO. Chihuahua: Mount Mohinora, 1 Sep 1898, E.W. Nelson 4895 (GH!; Isolectotypes: GH!, US, US-photo at GH!). The lectotype and isolectotype sheets at GH apparently hold the upper portion and base of a single plant, respectively, perhaps mounted separately by mistake.

Plants perennial. Stems erect, 18-40 cm tall, sometimes few branched from the base, invested with an understorey of minute, eglandular, spreading hairs, the overstorey absent or nearly so to much longer, stiffly spreading, vitreous hairs, the latter often gland tipped. Leaves sparsely and minutely scabridulous to sparsely glandular pilose, oblanceolate or narrowly oblong lanceolate to elliptic, 18-62 mm long, 5-13 mm wide, clasping to subclasping, sometimes barely so, 3(-5) veined, the veins strongly raised on the abaxial surface, margins of lower leaves sometimes with short, spreading cilia. Floral bracts entire, 3 veined, 5-8 mm wide, the uppermost red tipped. Calyces red and glandular at the apex, 18-25 mm long, tubular or narrowing toward the apex, the primary lobes 5-9 mm long, with acute apices, nearly equal in length, the secondary lobes 2-4 mm long, triangular. Corollas 20-32 mm long, the lower lip of 3, thick, green teeth ca. 2 mm long, the galea 8-12 mm long, pilose and glandular dorsally, ca. 33%-40% as long as the corolla, exerted 2-10 mm from the calyces. Chromosome number unknown.

Northern Chihuahua to central Durango (Map 2); pine-oak to pine woodlands, 2100-2670 m; Jul-Sep.

Additional collections examined: MEXICO. Chihuahua: Mpio. Bocoyna, W of Creel on mesa and E slope above Rio Oteros, 1 Aug 1977, Bye & Weber 7829 (GH); near Colonia Garcia, 25 Aug 1899, Nelson 6101 (GH); S of Colonia Garcia, 23 Sep 1934, Pennell 19181 (GH); Mpio. Guerrero, 7 mi E of Tomochic on road to La Junta and Cuauhtemoc, 5 Oct 1986, Spellenberg et al. 8880 (TEX); near Colonia Garcia, 4 Jun 1899, Townsend & Barber 8 (GH, MO); near

Colonia Garcia, 9 Aug 1899, Townsend & Barber 250 (GH, MO). Durango: Mpio. Santiago Papasquiaro, ca. 22 air km WNW of Santiago Papasquiaro, 25 Aug 1983, Worthington 11437 (NY).

The vestiture of *Castilleja aspera* appears to be somewhat more variable than normal for most species of the genus. The type of *C. nelsonii* represents an extreme condition with strongly reduced vestiture, the leaves only sparsely and minutely scabrid. Worthington 11437 from northern Durango is similar to the type of *C. aspera* and other plants from around Colonia Garcia, Chihuahua, except for Pennell 19181 and Bye & Weber 7829, which are more like the type of *C. nelsonii*. Additionally, the secondary calyx lobes of *C. nelsonii* (the type) are well differentiated but not sharply acute as in the other specimens of *C. aspera*, and it was apparently solely on this basis that Eastwood separated *C. nelsonii* from *C. aspera*. Nevertheless, considering their morphological similarity in all other features, as well as their geographical identity, the two are treated here as conspecific.

Plants from southcentral Durango have been identified by Eastwood (1909) and others as *Castilleja aspera*, but in the present study, they primarily are placed with following species, which is previously undescribed.

Castilleja durangensis Nesom, sp. nov. TYPE: MEXICO. Durango: ca. 5 km NE of El Salto, Rte 40 near K 1059, pine woods and wet llanos, under pine trees, with *Eryngium*, *Commelina*, *Ranunculus*, *Calochortus*, and *Lobelia*, 2780 m, 6 Aug 1966, R.W. Cruden 1160, voucher for chromosome count of $n=12$ [Heckard 1968, as *Castilleja aspera*] (HOLOTYPE: TEX!; Iso-type: GH!)

Castilleja asperae Eastwood similis sed vestimento eglanduloso dense brevithispidulo et foliis angustioribus absque venis valde elevatis differt.

Perennials, apparently from shallow, slender, woody rhizomes; stems, leaves, and bracts evenly and densely hispidulous with sharp pointed, spreading, sometimes slightly deflexed hairs 0.1-0.3(-0.5) mm long, eglandular. Stems erect, unbranched, 13-40 cm tall. Leaves ascending, often distinctly recurved, linear to oblong lanceolate, basally rounded and subclasping, 2-5 cm long, 2-5 mm wide, 3(-5) veined. Mature inflorescence/inflorescence (3-)-6-15 cm long; floral bracts entire, narrowly oblanceolate to narrowly obovate, 3 veined, 5-8 mm wide, red tipped. Calyces red and glandular at the apex, 18-22 mm long, tubular, the primary lobes 6-7 mm long, with acute apices, nearly equal in length, the secondary lobes 1-4 mm long, triangular. Corollas 21-25 mm long, the lower lip of 3, thick, green teeth 1-2 mm long, the galea 7-10 mm long, pilose and glandular dorsally, ca. 33%-40% as long as the corolla, exerted 2-6 mm from the calyces. Chromosome number, $n=12$ (voucher is type collection).

Southern Durango (Map 2); grassy slopes, sometimes wet, in pine and pine-oak woodlands; (2300)-2500-2850 m; Jun-Sep(-Oct).

Additional collections examined: MEXICO. Durango: 32.7 mi W of Durango, well spaced pine-oak forest on gentle slopes of loamy soils over volcanic rock, 23 Jul 1935, *Johnston 2686* (TEX); 39 mi E of El Salto, 7 Jun 1967, *Moldenke 1573* (NY, mixed with *Castilleja scorzoneraefolia*); 4 mi E of El Salto on road to Durango, 22 Aug 1957, *Ornduff & Solbrig 4639* (GH); Otinapa, 25 Jul-5 Aug 1906, *Palmer 367* (GH, MO, cited by Eastwood as *C. aspera*); El Salto (Aserraderos), grassy pinelands, 28 Aug 1934, *Pennell 18294* (GH, with 1 plant of *C. nervata*); Metates, N of Cuerva, pinelands on mt. slope, 29-30 Aug 1934, *Pennell 18993* (GH, with 1 plant of *C. nervata*); El Salto (Aserraderos), grassy pinelands, 31 Aug 1934, *Pennell 18499* (GH); El Salto (Aserraderos), mossy, grassy pinelands, 1 Sep 1934, *Pennell 18542* (GH); El Salto (Aserraderos), marshy glade in pinelands, 1 Sep 1934, *Pennell 18549* (GH); 3 mi E of El Salto off Hwy 40, 21 Sep 1974, *Rollins & Roby 7426* (GH, with 1 plant of *C. salensis* Eastwood, NY); 28 mi E of El Salto, 23 Jul 1976, *Walker 76H29* (MO).

The plants of *Castilleja durangensis* are confined to a small area of southern Durango (Map 2). They are somewhat similar in habit to *C. scorzoneraefolia*, but their evenly and densely short hispidulous vestiture, lacking glandular hairs, separates them from all other Mexican species. The secondary calyx lobes of *C. durangensis* are triangular with acute apices, a distinctive feature shared with plants of *C. aspera* and perhaps indicating a close relationship between the two. Plants of *C. aspera* are generally taller and the leaves broader than in *C. durangensis* and typically produce a vestiture more similar to that of *C. scorzoneraefolia*. The vestiture of the two branches of *Moldenke 1573* (*C. durangensis*) is slightly glandular, probably reflecting the genetic influence of nearby *C. scorzoneraefolia*, plants of which are mounted on the same sheet.

V. A new species related to *Castilleja lanata*

Castilleja galehintoniae Nesom, sp. nov. TYPE: MEXICO. Nuevo León. Mpio. Galeana, San José de Las Joyas, 2480 m, common on bare hillside, flowers yellow, 5 Jul 1983, *Hinton et al. 18422* (HOLOTYPE: TEX; Isotypes: ANSMI, MEXU, NY, UCJ).

Differs from *Castilleja lanata* A. Gray stature brevior, bracteis ac calycibus absque apicibus rubris, et corollis brevioribus labio infero loborum longorum linearium composito.

Perennials, arising from slender, woody, rhizome-like caudex branches, the stems 5-25 cm tall; stems, leaves, bracts, and calyces evenly and densely tomentose-villose with whitish, long, minutely filiform and intertwined hairs,


at least many of these originating as lateral branches of dendritic hairs, mostly with one node and several branches, minute glands sometimes apparent but never prominent beneath the other hairs. Leaves densely arranged on the stems, linear, 3 veined, 1.5-4.5 cm long, 1.0-1.5(-2.5) mm wide. Mature inflorescence/inflorescence 2-4 cm long; floral bracts 13-22 mm long, abruptly differentiated from the leaves, linear with a pair of linear, lateral lobes originating at about midlength. Calyces 15-18 mm long, tubular or slightly narrowing toward the apex, equally divided, the primary lobes 6-7 mm long, the secondary lobes rounded, barely developed, 0.5-1.0 mm long. Corollas 15-20 mm long, yellow with blackish veins, the lower lip of 3, thick, linear lobes 2-4 mm long, the galea 10-11 mm long, 50%-55% as long as the corolla, dorsally glandular but without other hairs, exerted 1-4 mm from the calyx. Chromosome number unknown.

Nuevo León endemic (Map 3); apparently restricted to gypsaceous substrates, in matorral, grasslands, cedar savannas, open oak woods; 1300-2480 m; Apr-Aug(-Oct).


Additional collections examined. MEXICO. Nuevo León: Mpio. Aramburi, E of Aramburi, 14 May 1991, *Hinton et al. 20939* (TEX). Mpio. Galeana: above E Carrizo, 16 Oct 1983, *Hinton et al. 18149* (TEX); S slope of Potosí, 25 Jun 1983, *Hinton et al. 18449* (TEX, dup); San José Las Joyas, 23 Jul 1983, *Hinton et al. 18590* (TEX, dup); road to Dieciocho de Marzo, 11.5 mi E of jct with Hwy 57 at San Rafael, 28 mi NW of Dieciocho de Marzo, 4 Jul 1985, *Luckow 2651* (TEX); 15 mi SW of Galeana, above Encinal, 19 May 1934, *Mueller 465* (GH, TEX); ca. 15 mi SW of Galeana, Taray to Alamar at la Mesa de la Zorra, 20 Jul 1934, *Mueller 1109* (GH, TEX); Arroyo Hondo, Hacienda San José de Raices, 31 Jul 1935, *Mueller 2907* (GH, MO, TEX); Hacienda Pabillo, 1 Aug 1936, *Taylor 41* (MO, TEX); Hacienda Pabillo, 26 Aug 1936, *Taylor 221* (TEX).

The new species is endemic to southern Nuevo León, México (Map 3), and it is named for Gale Hinton, daughter of Jaime Hinton and brother of George, who loved the mountains and flowers around her home in the same area of Nuevo León. *Castilleja galehintoniae* is clearly most closely related to *C. lanata*, which also produces a dense vestiture of branched hairs, mostly linear leaves, an evenly divided calyx, and corollas with a lower lip of 3 linear lobes. The two species differ in features noted in the following couplet.

1. Stems 20-100 cm tall; floral bracts with lobes usually originating from the proximal third, densely lanate basally, the apices glandular but not densely lanate; calyces 20-27 mm long, with primary lobes 10-14 mm long; corollas 22-37 mm long, the galea dorsally pilose as well as glandular. *C. lanata*
1. Stems 5-25 cm tall; floral bracts with lobes usually originating from near the middle, apices densely lanate, the vestiture similar from base to tip;



Map 3. Distribution of *Castilleja galehintoniae* and *C. lanata* in Nuevo León and closely adjacent areas. Arrow shows the location of the type collection of *C. galehintoniae*; the type of *C. lanata* is from Texas.



Map 4. Distribution of *Castilleja bella*. Arrow shows the location of the type collection.

calyces 15-18 mm long; with primary lobes 6-7 mm long; corollas 15-20 mm long; the galea dorsally glandular but not otherwise pubescent. *C. galehintoniae*

While *Castilleja lanata* is somewhat variable in size, it is consistently and easily recognizable as a single species over its entire geographic range. It occurs widely in the southwestern United States (Arizona, Colorado, New Mexico, and southwest Texas) and northern México (from Sonora, Chihuahua, and northern Durango to Coahuila, northern Zacatecas, San Luis Potosí, and southern Tamaulipas). It is also relatively common in western Nuevo León, where its geographic range touches that of *C. galehintoniae*, and the two are apparently even slightly sympatric in that area (Map 3). Several collections from the area of sympatry can be identified as possible intermediates (e.g., Cowan 4626-TEX, near the village of El Potosí, and some plants of Hinton 18449-identified here as *C. galehintoniae*), but where the two taxa meet, the transition in morphology for the most part appears to be abrupt, with *C. galehintoniae* strikingly different in appearance from *C. lanata*. Putative intermediates have shorter calyces and corollas than normal for *C. lanata*, but the floral bracts and calyces are distally distinctly red with somewhat reduced vestiture. Some other collections of *C. lanata* from this area also are unusual in producing lobed upper leaves, although the origin of this variability is not clear.

VI. A new species of uncertain relationships

Castilleja dendridion Nesom, sp. nov. TYPE: MEXICO, Oaxaca: Dist. Centro, 10 mi NE of Oaxaca along Hwy 175, pine-oak forest on steep slopes, 2 Sep 1982, L.E. Gieschen s.n. (HOLOTYPE: TEX¹).

A speciesbus ceteris *Castillejæ* dignoscenda vestimento denso trichomatium dendriticorum flavipigmentosorum, foliis confertim dispositis linearibus integrisque, bracteis floralibus lobis ad apicem expansis dentatisque, et lobis calycum pariter divisis.

Shrubby perennials, the stems basally woody and at least 40 cm tall; stems, leaves, floral bracts, and calyces densely invested with multicellular, dendritic hairs 0.2-1.0 mm high, branching 2-4 times above the 1-2 basal (stipe) cells, each hair with 2(-3) branches at each cellular node, the hairs of the stems and leaves usually prominently yellow pigmented, giving the whole plant a yellowish cast. Leaves linear-filiform, 9-17 mm long, ca. 1 mm wide, not at all basally widened or clasping, densely arranged (3-6 per cm of the stem) with axillary tufts of smaller leaves at nearly every node. Mature inflorescence/inflorescence 4-10 cm long. Floral bracts abruptly differentiated from

the leaves, 14-15 mm long, 2-3 mm wide at the base, with a pair of linear-lanceolate lobes arising 2-3 mm from the base, all 3 lobes red and abruptly broadened apically, the apices truncate and shallowly toothed. Calyces red tipped and glandular, 13-15 mm long, tubular, equally divided, the primary lobes 6-8 mm long, secondary lobes delatate, ca. 1 mm long. Corollas 14-15 mm long, lower lip of 3, thick teeth ca. 2 mm long; the galea 8-9 mm long, ca. 60% as long as the corolla, dorsally glandular but without other hairs, exerted (0-)1-2 mm from the calyx. Chromosome number unknown.

Known only from the type collection (Map 1), which comprises a single, though full and densely floriferous, branch.

Castilleja dendridion is so distinctive in its morphology that no other species in the genus can be easily identified as closely related. Other species of *Castilleja* produce dendritic hairs (see Nesom 1991, as well as *C. galehintoniae*, above), but such highly elaborated hairs are not found elsewhere in the genus. Further, the peculiar morphology of the floral bracts apparently is found only in this species. The evenly divided calyx at least suggests that *C. dendridion* belongs among the species placed by Eastwood (1909) in sect. *Euchroma* (Nutt.) Benth., although the overview of the subtribe *Castillejinæ* by Chuang & Heckard (1991) suggests that this morphology is probably primitive within the subtribe. Chuang & Heckard (1991) noted that overemphasis of the nature of calyx incision in classification probably results in the recognition of artificial groups, and they observed that satisfactory arrangement of *Castilleja* species into groups awaits a more detailed knowledge of the genus. All taxa treated in the present study are in *Castilleja* subg. *Castilleja* sensu Chuang & Heckard.

ACKNOWLEDGMENTS

I thank Dr. T.I. Chuang and Dr. B.L. Turner for their comments on the manuscript and the staffs of GH, NY, and PH for loans of specimens. Specimens from MO were studied on a recent visit there.

LITERATURE CITED

- Brandegee, T.S. 1914. *Plantae Mexicanæ Purpusianæ*, VI. Univ. Calif. Publ. Bot. 6:51-77.
- Breedlove, D.E. & L.R. Heckard. 1970. *Gentrya*, a new genus of Scrophulariaceae from Mexico. *Brittonia* 22:20-24.
- Chuang, T.I. & L.R. Heckard. 1991. Generic realignment and synopsis of Subtribe *Castillejinæ* (Scrophulariaceae - Tribe Pedicularae). *Syst. Bot.* 16:644-666.

- Crosswhite, F.S. 1970. *Castilleja roei* (Scrophulariaceae). Amer. Midl. Nat-uralist 83:630-631.
- Eastwood, A. 1909. Synopsis of the Mexican and Central American species of *Castilleja*. Contr. Gray Herb., n. ser. 36:563-591.
- Heckard, L.R. 1968. Chromosome numbers and polyploidy in *Castilleja* (Scrophulariaceae). Brittonia 20:212-226.
- Holmgren, N.H. 1970. *Castilleja*. Pp. 1439-1442 in Correll, D.S. & M.C. Johnston. *Manual of the Vascular Plants of Texas*. Texas Research Foundation, Renner, Texas.
- _____. 1976. Four new species of Mexican *Castilleja* (subgenus *Castilleja*, Scrophulariaceae) and their relatives. Brittonia 28:195-208.
- _____. 1978. *Castilleja* (Scrophulariaceae) of Costa Rica and Panama. Brittonia 30:182-194.
- Jiménez, C.R. 1985. *Castilleja*. Pp. 346-349, in Rzedowski, J. & G.C. Rzedowski (eds.). *Flora Fanerogámica del Valle de México*. Vol. II. Dicotyledoneae. Instituto de Ecología, México, D.F.
- Levin, G.A. & R. Moran. 1989. The vascular flora of Isla Socorro, Mexico. Mem. San Diego Soc. Nat. Hist. 16:1-71.
- McDonald, J.A. 1990. The alpine-subalpine flora of northeastern Mexico. Sida 14:21-28.
- _____. In press. Phytogeography of the alpine-subalpine flora of north-eastern Mexico. In T.P. Ramamoorthy, R. Bye, A. Lot, & J. Fa (eds.), *Biological Diversity of Mexico: Origins and Distribution*. Oxford Press, New York, New York.
- Nesom, G.L. 1992. A new species of *Castilleja* (Scrophulariaceae) from southcentral Texas, with comments on other Texas taxa. *Phytologia* 72:209-230.
- Rzedowski, J. 1975. Tres dicotiledoneas nuevas de interes ornamental. *Bol. Soc. Bot. México* 35:40-47.
- Standley, P.C. 1936. Studies of American plants - VI. Field Mus. Publ. Bot. 11:145-276.
- _____. 1940. Studies of American plants - X. Field Mus. Pub. Bot. 22:65-129.
- Standley, P.C. & J.A. Steyermark. 1944. Studies of Central American plants - IV. Publ. Field Mus. Nat. Hist. Bot. Ser. 23:31-109.